ORTHOPAEDIC ONCOLOGY · PATIENT GUIDE

Sarcoma Found After Lump Removal: Why Another Operation May Be Needed

A guide to specialist review, further investigations and the decisions that follow.

Dr. Vishnu Baburaj

Published 3 November 2026 · Updated 3 November 2026 · Clinically reviewed 3 November 2026

https://drvishnubaburaj.com/blog/oncology/sarcoma-found-after-lump-removal/



You may have expected the lump removal to finish the matter. Then the tissue report says “sarcoma”. When a sarcoma was removed without the investigations and surgical planning needed for that diagnosis, clinicians call it an *unplanned excision*.[1]

This article is for someone who already has that unexpected diagnosis, particularly after removal of a lump in a limb or the trunk. The first discussion can focus on what is known now and what needs to happen next.

Start with a specialist review

A sarcoma team coordinates specialist assessment and treatment, drawing on surgical, imaging, pathology and oncology expertise.[7] The doctor who received the result should help arrange that referral and a contact for updates.

For a pathology second opinion, the reviewing pathologist may need the original slides or stored tissue blocks.[3]

The review needs to establish what was removed, whether the specimen was intact or fragmented, and what the margins showed. Those details help the team interpret the scan and assess whether tumour may remain.[6]

Useful records include the operation note, pathology report, discharge summary and any scans taken before removal. You can share what you already have and where the earlier care took place; the team should help obtain missing material.

> **Dr. Vishnu's perspective**
>
> An unexpected report can leave you wondering what the first operation means for everything that follows. I want to review the information with you before drawing conclusions. If another operation is being considered, I would explain why, without jumping to conclusions about blame before the original records have been reviewed. We should leave that discussion knowing the next step.

What the scans are trying to find out

> Two separate questions after removal

Assessment	What it asks
The operated area	Is there visible tumour remaining, and what area needs surgical planning? MRI can help.[5]
Staging	What is the extent of the cancer, including whether it has spread elsewhere?[4]

After an unplanned excision, swelling and fluid from the operation can make MRI harder to interpret. The scan needs to be read alongside the operation details and pathology.[6]

An MRI showing no visible tumour cannot reliably exclude remaining sarcoma cells. It helps planning, but cannot give a microscopic “all clear”. That is why a further operation may still be discussed when there is no lump left to feel.[5][2]

More about the evidence

A small retrospective study compared MRI with findings at a second operation and found that MRI missed some remaining disease.[5] A separate retrospective referral-centre study found some scans equivocal and stressed interpretation with the operation and pathology.[6]

Why another operation may be advised

The aim of *re-excision* is to remove the previous surgical area with an appropriate margin. This can involve more tissue than the visible scar because the first procedure may have exposed surrounding tissues to tumour. Skin grafts or other reconstructive procedures may be needed to close the resulting defect.[1]

Current GEIS guidance recommends considering re-excision after an unplanned, inadequate removal. The decision belongs within a specialist team discussion.[7] Ask the surgeon to show you the proposed area of removal and explain the expected effect on movement, work and daily activities.

Does everyone need further surgery?

Re-excision may be advised when adequate clearance is achievable with acceptable harm. If another operation would cause considerable harm or is unlikely to clear the affected area, UK guidance allows consideration of radiotherapy or observation instead.[2]

Radiotherapy is a separate decision. Its role depends on the tumour and the overall treatment plan; it is not an automatic replacement for obtaining adequate surgical margins when further surgery is feasible.[7]

> **Ask for the reasoning:** "What makes re-excision suitable for me, what would it involve, and what are the alternatives?" If observation is proposed, ask for the written follow-up plan and the person responsible for reviewing it.

What can the research tell you about the outlook?

Research links unplanned excision and residual tumour with poorer control of sarcoma in the operated area. This supports careful assessment after an unexpected diagnosis. It does not establish that the first operation caused a particular person's later outcome or provide a personal prediction.[8]

More about the evidence

The systematic review combined observational studies. The groups differed in factors such as tumour size and depth. An association between unplanned excision and local recurrence cannot, by itself, establish cause and effect; the diagnosis and subsequent treatment also matter.[8]

Agree the next steps before you leave

Research has not established one ideal interval between unplanned removal and re-excision.[1] The team should coordinate the timetable and explain which results are still needed before the next decision.

- Has the diagnosis been reviewed by a sarcoma pathologist?
- What do the first operation and margin findings tell us?

- Which scans are needed, and what will each answer?
- Is more surgery recommended, and what would be removed?
- Would reconstruction or radiotherapy be part of the plan?
- Who should I contact if an appointment or result is delayed?

A brief written plan can make the next step easier to follow, including the next appointment and a contact if something is delayed.

Frequently Asked Questions

Why discuss more surgery when the lump has already gone?

Microscopic tumour may remain after an unplanned removal. Re-excision aims to clear the previous surgical area. Whether it is appropriate needs specialist assessment. [6][2]

Does an MRI showing no tumour mean no further treatment is needed?

No. MRI helps with planning, but it cannot reliably exclude residual sarcoma after unplanned excision. The team also considers the pathology and the first operation. [5][6]

Does everyone need another operation after an unplanned sarcoma removal?

No. If further surgery would cause considerable harm or is unlikely to clear the affected area, a specialist team may consider radiotherapy or observation instead. These are individual decisions. [2]

How soon should a second operation happen?

Research has not established one suitable interval for everyone. Arrange specialist review promptly and ask for a coordinated timetable for the investigations and treatment recommended for you. [1]

Your consultation

Plan a review after unexpected sarcoma findings in Kannur

After a sarcoma diagnosis following lump removal, you can book a consultation with Dr. Vishnu Baburaj at Aster MIMS Kannur to discuss the findings and proposed next steps.

Book a consultation Plan your visit

Travelling to Kannur

For a visit from outside Kannur, the booking discussion can cover the appointment and which existing records are needed.

Preparing for your consultation

> For referring clinicians: enquire about arranging an assessment through the existing appointment contacts.

The evidence behind this article

1. Nakamura T, Hasegawa M. Unplanned Excision in Soft Tissue Sarcoma: Current Knowledge and Remaining Gaps. *Diagnostics*. 2025;15:453.
2. Hayes AJ, Nixon IF, Strauss DC, et al. UK guidelines for the management of soft tissue sarcomas. *British Journal of Cancer*. 2025;132:11–31. Published online in 2024.
3. National Cancer Institute. Surgical Pathology Reports. Reviewed August 2022.
4. National Cancer Institute. Cancer Staging.
5. Alramdan MHA, Kasalak Ö, Been LB, et al. MRI after Whoops procedure: diagnostic value for residual sarcoma and predictive value for an incomplete second resection. *Skeletal Radiology*. 2021;50:2213–2220.
6. Gingrich AA, Elias A, Lee CYM, et al. Predictors of Residual Disease after Unplanned Excision of Soft Tissue Sarcomas. *Journal of Surgical Research*. 2017;208:26–32. Published online in 2016.
7. Vaz-Salgado MA, Valverde-Morales C, Alvarez R, et al. Diagnosis and Therapy of Soft Tissue Sarcomas: Spanish Group for Research in Sarcomas (GEIS) Guidelines. *Cancers*. 2025;17:3158.
8. Larios F, Gonzalez MR, Ruiz-Arellanos K, et al. Is Unplanned Excision of Soft Tissue Sarcomas Associated with Worse Oncological Outcomes? A systematic review and meta-analysis. *Cancers*. 2024;16:443.

Medically reviewed by Dr. Vishnu Baburaj, MS Ortho (PGI Chandigarh), MRCS (UK) · Last reviewed Nov 3, 2026

This article is for educational purposes and does not replace a medical consultation. For personalised advice, book an appointment at Aster MIMS Kannur.